

**AN AXIOMATIC CHARACTERIZATION OF THE DIMENSION  
OF SUBSETS OF EUCLIDEAN SPACES**

Yoshiaki HAYASHI

*Institute of Mathematics, Konan University, Higashinada, Kobe 658, Japan*

Received 7 October 1988

Revised 26 July 1989

Menger's axioms on dimension functions are not adequate to determine the dimension on the class of all subspaces of Euclidean spaces. In this paper, the following is shown: A decomposition axiom together with Menger's axioms characterize the dimension on the class of all subspaces of Euclidean spaces, and the axioms are independent.

AMS (MOS) Subj. Class: 54F45

dimension      axiomatic characterization      Menger's axioms

**1. Axioms**

Let  $\mathcal{Q}$  be a class of spaces,  $\mathcal{E}$  the class of all subspaces of Euclidean spaces and  $\mathcal{S}$  the class of all separable metric spaces. In 1929, Menger studied real-valued set functions  $d$  defined on the class  $\mathcal{Q}$  satisfying the following conditions (A1)–(A5):

- (A1) monotone: if  $X \in \mathcal{Q}$  and  $X' \subset X$ , then  $d(X') \leq d(X)$ ,
- (A2)  $F_\sigma$ -constant: if  $X (\in \mathcal{Q})$  is the countable sum of closed sets  $X_i$ , then  $d(X) \leq \max_i d(X_i)$ ,
- (A3) topological: if  $X (\in \mathcal{Q})$  and  $X'$  are homeomorphic, then  $d(X) = d(X')$ ,
- (A4) compactifiable: each  $X (\in \mathcal{Q})$  is homeomorphic to a subset of a compact space  $A (\in \mathcal{Q})$  such that  $d(X) = d(A)$ ,
- (A5) normed:  $d(\emptyset) = -1$ ,  $d(I^n) = n$  ( $n = 0, 1, 2, \dots$ ) for the  $n$ -dimensional interval  $I^n$ .

Menger raised a problem: “In the case  $\mathcal{Q} = \mathcal{E}$ , does  $d$  coincide with the dimension  $\dim$ ?”, and he proved that if  $\mathcal{Q}$  is the class of all subspaces of the 2-dimensional Euclidean space, then  $d$  coincides with  $\dim$  [7].

In 1968, Shvedov showed that the conditions (A1)–(A5) are not adequate to determine the dimension in the case  $\mathcal{Q} = \mathcal{E}$  (this result is introduced in [4]).

In order to characterize dimension, what kind of conditions should be added besides the above conditions (A1)–(A5)? Many proposals regarding characterization of dimension axiomatically are already known (for instance, [1, 3, 6, 8–13] (for metric spaces)). However, as far as we know, there has been no proposal complementing the idea of Menger. In the present paper, we prove that the condition

(A6) if  $d(X) < n$  (where  $n$  is a positive integer), then there exist  $n$  sets  $X_i$  such that  $X = \bigcup_{i=1}^n X_i$  and  $d(X_i) \leq 0$  ( $i = 1, 2, \dots, n$ ) together with the conditions (A1)–(A5) characterize dimension in the case  $\mathcal{Q} = \mathcal{E}$ , and that the conditions (A1)–(A6) are independent.

Throughout this paper, spaces are subspaces of Euclidean spaces unless otherwise specified.

In cases  $\mathcal{Q} = \mathcal{E}$  or  $\mathcal{Q} = \mathcal{S}$ , as  $\dim$  is a function satisfying the conditions (A1)–(A6), it is obvious that the conditions (A1)–(A6) are consistent and that in order that a set function  $d$  defined on  $\mathcal{Q}$  coincides with  $\dim$  it is necessary that  $d$  satisfies the conditions (A1)–(A6).

## 2. Definitions

Throughout this paper,  $I$  is the unit closed interval of real numbers, i.e.,  $I = \{x \mid x \text{ is a real number, } 0 \leq x \leq 1\}$ ,  $R^m$  is the  $m$ -dimensional Euclidean space, and  $I^m$  is the unit cube in  $R^m$ . The set of all points  $x \in I^m$  such that at most  $n$  of the coordinates of  $x$  are irrational is denoted by  $I_n^m$ . Each point  $x$  in  $R^m$  is represented by  $(x^{(1)}, x^{(2)}, \dots, x^{(m)})$ , where  $x^{(i)}$  is the  $i$ th coordinate of  $x$ . A *parallelotope* in  $R^m$  is a set having the form  $\{x \in R^m \mid a_i \leq x^{(i)} \leq b_i \text{ (} i = 1, 2, \dots, m)\}$ , where  $a_i$  and  $b_i$  are real numbers with  $a_i < b_i$  ( $i = 1, 2, \dots, m$ ), and a *half open parallelotope* in  $R^m$  is a set having the form  $\{x \in R^m \mid a_i \leq x^{(i)} \leq b_i \text{ (} i = i[1], i[2], \dots, i[n]), a_i < x^{(i)} < b_i \text{ (} i \neq i[1], i[2], \dots, i[n])\}$ , where  $\{i[1], i[2], \dots, i[n]\}$  is a proper subset of  $\{1, 2, \dots, m\}$ . The closure of a half open parallelotope is a parallelotope. The definition of a *face* of a parallelotope should be evident. Each face of the closure of a half open parallelotope  $P$  is called a *face* of  $P$ . In  $R^m$ , an *n-parallelotope* ( $n < m$ ) is a set having the form  $\{x \in R^m \mid a_i \leq x^{(i)} \leq b_i \text{ (} i = i[1], i[2], \dots, i[n]), x^{(i)} = a_i \text{ (} i \neq i[1], i[2], \dots, i[n])\}$ , where  $\{i[1], i[2], \dots, i[n]\}$  is a proper subset of  $\{1, 2, \dots, m\}$ .

For any parallelotope  $P$ , we use  $g(P)$  to denote the barycenter (in the sense of Euclidean geometry) of  $P$ , and, for any half open parallelotope  $P'$ ,  $g(P')$  shall denote the barycenter of the closure  $\overline{P'}$  of  $P'$ .

The sum of finitely many parallelotopes is called a *polyparallelotope*, and the sum of finitely many half open parallelotopes is called a *half open polyparallelotope*. In  $R^m$ , if the intersection  $M$  of an  $(m-1)$ -dimensional hyperplane and the boundary (in the topological sense)  $\partial P$  of a polyparallelotope  $P$  contains an  $(m-1)$ -dimensional rectangle, then the intersection  $M$  is called a *face* of  $P$ , and each face of the closure  $\overline{P'}$  of a half open polyparallelotope  $P'$  is called a *face* of  $P'$ .

## 3. Compactification of $I_n^{2n+1}$

**Theorem 3.1.** *Let  $G$  be a  $G_\delta$ -set of  $I_n^{2n+1}$  ( $\subset R^{2n+1}$ ) containing  $I_n^{2n+1}$  and  $X$  be an  $n$ -dimensional space. Then  $G$  contains a subset which is homeomorphic to  $X$ .*

**Proof.** Let  $F = I^{2n+1} - G$ . Then  $F$  is an  $F_\sigma$ -set in  $I^{2n+1}$ . We may write  $F = \bigcup_{i=1}^{\infty} F_i$  (each  $F_i$  is a closed set of  $I^{2n+1}$ ). For each integer  $k$  ( $\geq 0$ ), we define the set of parallelotopes  $S(k)$  by induction as follows:

$S(0) = \{I^{2n+1}\}$ . Define  $S(k+1)$  ( $k \geq 0$ ), assuming that  $S(k)$  is defined, as follows:

$$S(k) = \{T(k, 1), T(k, 2), \dots, T(k, s(k))\}$$

is a set of parallelotopes in  $R^{2n+1}$ . Let  $T(k, i)$  be an element of  $S(k)$ ,  $a$  be a vertex of  $T(k, i)$ , and  $\{j[1], j[2], \dots, j[n+1]\} \subset \{1, 2, \dots, 2n+1\}$ . Let

$$\begin{aligned} P(k, i, a; j[1], [2], \dots, j[n+1]) \\ = \{x \in T(k, i) \mid x^{(j[l])} = a^{(j[l])} \ (l = 1, 2, \dots, n+1)\}. \end{aligned}$$

We write  $P(k, i, a; j[1], \dots, j[n+1])$  as  $P$  for short. Now assume that  $S(k)$  is defined so that every  $P$  is contained in  $I_n^{2n+1}$ .  $P$  does not intersect  $F_{k+1}$ . There is a rational number  $d(k)$  ( $> 0$ ) such that the parallelotopes

$$\begin{aligned} T(k, i, a; j[1], \dots, j[n+1]) \\ = \{x \in T(k, i) \mid a^{(j[l])} - d(k) \leq x^{(j[l])} \leq a^{(j[l])} + d(k) \ (l = 1, 2, \dots, n+1)\} \end{aligned}$$

do not intersect  $F_{k+1}$  for all combinations  $\{i, a, j[1], \dots, j[n+1]\}$  ( $i = 1, 2, \dots, s(k)$ ;  $a$  is some vertex of  $T(k, i)$ ;  $\{j[1], \dots, j[n+1]\} \subset \{1, 2, \dots, 2n+1\}$ ). Denote these parallelotopes  $T(k, i, a; j[1], \dots, j[n+1])$  by  $T(k, i)_1, T(k, i)_2, \dots, T(k, i)_t$ . Let

$$\begin{aligned} J(k, i; \alpha(1), \alpha(2), \dots, \alpha(p)) \\ = \left\{ x \mid x \in \bigcap_{q=1}^p T(k, i)_{\alpha(q)}, \right. \\ \left. x \notin T(k, i)_\gamma \text{ for each } \gamma \ (\gamma \neq \alpha(1), \alpha(2), \dots, \alpha(p)) \right\}. \end{aligned}$$

The closure of each  $J(k, i; \alpha(1), \dots, \alpha(p))$  is a parallelotope or  $\emptyset$ . Denote these parallelotopes by  $J(k, i; 1), J(k, i; 2), \dots, J(k, i; h(k))$ . Subdivide each  $J(k, i; h)$  into  $2^{2n+1}$  parallelotopes by means of  $2n+1$  hyperplanes which contain  $g(J(k, i; h))$  and are parallel to  $2n+1$  coordinate hyperplanes respectively (where  $g(J)$  is the barycenter of  $J$ ). Let  $S(k+1)$  be the set of all such parallelotopes produced from all elements of  $S(k)$ . Let

$$\begin{aligned} C(k) &= \{x \mid x \in T \text{ for some element } T \text{ of } S(k)\} \\ (k &= 1, 2, \dots). \end{aligned} \tag{1}$$

Then,

$$C(1) \supset C(2) \supset \dots, \quad \text{Mesh } S(k) \rightarrow 0 \quad (k \rightarrow \infty).$$

Now, let  $X$  ( $\in \mathcal{E}$ ) be an  $n$ -dimensional space. Since  $I^{2n+1} - I_n^{2n+1}$  is an  $n$ -dimensional universal set, there exists a compactification  $Y$  of  $X$  such that  $\dim X = \dim Y = n$  and  $Y \subset I^{2n+1} - I_n^{2n+1}$ . We define countably many half open polyparallelotopes  $K(k)$  ( $k = 1, 2, \dots$ ) as follows:

Let  $\{j[1], j[2], \dots, j[n+1]\} \subset \{1, 2, \dots, 2n+1\}$  in  $I^{2n+1}$ , and let

$$\begin{aligned} &L(j[1], j[2], \dots, j[n+1]) \\ &= \{x \in I^{2n+1} \mid x^{(j[1])} = x^{(j[2])} = \dots = x^{(j[n+1])} = \frac{1}{2}\} \end{aligned}$$

(we denote  $L(j[1], \dots, j[n+1])$  by  $L$  for short). It follows that  $L \cap Y = \emptyset$ . There exists a rational number  $r(1)$  such that the half open parallelotopes

$$\begin{aligned} &M(j[1], \dots, j[n+1]) \\ &= \{x \in I^{2n+1} \mid \frac{1}{2} - r(1) < x^{(j[l])} < \frac{1}{2} + r(1) \ (l = 1, 2, \dots, n+1)\} \end{aligned}$$

do not intersect  $Y$  for all combinations  $\{j[1], j[2], \dots, j[n+1]\} (\subset \{1, 2, \dots, 2n+1\})$ . Let

$$K(1) = \bigcup \{M(j[1], \dots, j[n+1]) \mid \{j[1], \dots, j[n+1]\} \subset \{1, 2, \dots, 2n+1\}\}.$$

Divide  $I^{2n+1} - K(1)$  into parallelotopes by means of all hyperplanes (of  $R^{2n+1}$ ) each of which contains some face of  $K(1)$ , and let  $A(1)$  be the set of all these parallelotopes produced in this way. Subdivide each  $V (\in A(1))$  into parallelotopes by means of  $2n+1$  hyperplanes which contain  $g(V)$  and are parallel to  $2n+1$  coordinate hyperplanes respectively. Let  $B(1)$  be the set of all parallelotopes produced from all elements of  $A(1)$  in this way. Assume that the half open polyparallelotope  $K(k)$  is defined and

$$K(k) = \bigcup \{M_\beta \mid \beta = 1, 2, \dots, \zeta(k)\}$$

where each  $M_\beta$  is a maximal half open parallelotope contained in  $K(k)$ . We shall define  $K(k+1)$  using  $K(k)$ . Let  $H$  be the sum of all hyperplanes (in  $R^{2n+1}$ ) each of which contains some face of  $K(k)$ .  $H$  divides  $I^{2n+1}$  into a finite number  $u$  of parallelotopes  $N_1, N_2, \dots, N_u$ . For any  $N_i$  ( $1 \leq i \leq u$ ), either  $N_i$  is contained in the closure  $\overline{M_\beta}$  (in  $R^{2n+1}$ ) of some  $M_\beta$  or  $N_i$  does not contain any inner-point of  $M_\beta$  for all  $\beta$ . Let  $N'_1, N'_2, \dots, N'_{u'}$  be those  $N_i$  which do not contain any inner-point of  $M_\beta$  for all  $\beta$ . For each  $N'_s$  ( $1 \leq s \leq u'$ ), subdivide  $N'_s$  into parallelotopes by means of  $2n+1$  hyperplanes which contain  $g(N'_s)$  and are parallel to  $2n+1$  coordinate hyperplanes respectively. Let  $B(k) = \{W_1, W_2, \dots, W_\eta\}$  be the set of all such parallelotopes produced from  $N'_1, \dots, N'_{u'}$  in this way. Let  $\{j[1], j[2], \dots, j[n+1]\} \subset \{1, 2, \dots, 2n+1\}$ . For each element  $W_\varepsilon$  of  $B(k)$ , we define

$$\begin{aligned} &L(W_\varepsilon; j[1], \dots, j[n+1]) \\ &= \{x \in W_\varepsilon \mid x^{(j[l])} = g(W_\varepsilon)^{(j[l])} \ (l = 1, 2, \dots, n+1)\} \end{aligned}$$

(we denote  $L(W_\xi; j[1], \dots, j[n+1])$  by  $L$  for short). Then,  $L \cap Y = \emptyset$ . There exists a rational number  $r(k+1)$  such that the half open parallelotopes

$$\begin{aligned} & M(W_\xi; j[1], \dots, j[n+1]) \\ &= \{x \in W_\xi \mid g(W_\xi)^{(j[l])} - r(k+1) < x^{(j[l])} < g(W_\xi)^{(j[l])} + r(k+1) \\ &\quad (l = 1, 2, \dots, n+1)\} \end{aligned}$$

do not intersect  $Y$  for all combinations  $\{W_\xi, j[1], \dots, j[n+1]\}$  ( $W_\xi \in B(k)$ ,  $\{j[1], \dots, j[n+1]\} \subset \{1, 2, \dots, 2n+1\}$ ). Let

$$\begin{aligned} K(k+1) = & \bigcup \{M(W_\xi; j[1], \dots, j[n+1]) \mid W_\xi \in B(k), \\ & \{j[1], \dots, j[n+1]\} \subset \{1, 2, \dots, 2n+1\}\}. \end{aligned}$$

Thus  $K(k)$ ,  $B(k)$  ( $k = 1, 2, \dots$ ) are defined.

$B(k)$  is similar to  $S(k)$ , in the sense of classes of (point-) sets, and, setting

$$\begin{aligned} D(k) = & \{x \mid x \in V \text{ for some } V \in B(k)\} \quad (k = 1, 2, \dots), \\ D(1) \supset & D(2) \supset \dots, \quad \text{Mesh } B(k) \rightarrow 0 \quad (k \rightarrow \infty). \end{aligned}$$

There is a bijection  $f$  which maps  $\bigcup_{k=1}^{\infty} B(k)$  onto  $\bigcup_{k=1}^{\infty} S(k)$  such that

(i)  $f$  maps  $B(k)$  onto  $S(k)$  and the correspondence  $V (\in B(k)) \leftrightarrow f(V) (\in S(k))$  is similar in the sense of classes of (point-) sets ( $k = 1, 2, \dots$ ),

(ii) if  $V \in B(k)$ ,  $V' \in B(k+1)$  and  $V \supset V'$ , then  $f(V) \supset f(V')$ .

Therefore, if we set  $K = \bigcup_{k=1}^{\infty} K(k)$ , then the complement  $I^{2n+1} - K$  of  $K$  is homeomorphic to  $C (= \bigcap_{k=1}^{\infty} C(k))$ . Since  $I^{2n+1} - K$  contains  $Y$ ,  $C$  has a subset which is homeomorphic to  $Y$ . Therefore, from  $C \subset G$ ,  $G$  has a subset which is homeomorphic to  $Y$ , and therefore has a subset which is homeomorphic to  $X$ .  $\square$

**Theorem 3.2.** Let  $C (\in \mathcal{E})$  be any compactification of  $I_n^{2n+1}$ . Then, for an arbitrary  $n$ -dimensional space  $X (\in \mathcal{E})$ ,  $C$  has a subset which is homeomorphic to  $X$ .

**Proof.**  $C \subset R^m$  for some integer  $m (> 0)$  because  $C \in \mathcal{E}$ . We may assume  $m \geq 2n+1$ . Let

$$\begin{aligned} I[n] = & \{x \in I^m \mid \text{at most } n \text{ elements of } \{x^{(1)}, x^{(2)}, \dots, x^{(2n+1)}\} \text{ are} \\ & \text{irrational, } x^{(l)} = 0 \ (l = 2n+2, 2n+3, \dots, m)\}. \end{aligned}$$

Then  $I[n]$  is homeomorphic to  $I_n^{2n+1}$  and  $C$  is a compactification of  $I[n]$ . Let  $I[n]'$  ( $\subset C$ ) be a set which is homeomorphic to  $I[n]$ , and let  $f$  be a homeomorphic mapping of  $I[n]$  onto  $I[n]'$ . There exist two  $G_\delta$ -sets  $G, G'$  (of  $R^m$ ) and a homeomorphic mapping  $f': G \rightarrow G'$  such that  $G$  and  $G'$  contain  $I[n]$  and  $I[n]'$ , respectively, and  $f'$  is an extension of  $f$  [5].  $(f')^{-1}(G' \cap C)$  is a  $G_\delta$ -set (of  $R^m$ ) containing  $I[n]$ . Let

$$\begin{aligned} E[n] = & ((f')^{-1}(G' \cap C)) \cap \{x \in R^m \mid 0 \leq x^{(l)} \leq 1 \ (l = 1, 2, \dots, 2n+1), \\ & x^{(l)} = 0 \ (l = 2n+2, 2n+3, \dots, m)\}, \end{aligned}$$

then  $E[n]$  is a  $G_\delta$ -set (of  $R^m$ ) containing  $I[n]$ . If we set

$$E[n]'' = \{y \in R^{2n+1} \mid \exists x (\in E[n]) \ y^{(l)} = x^{(l)} \ (l = 1, 2, \dots, 2n+1)\},$$

then  $E[n]''$  is a  $G_\delta$ -set (of  $R^{2n+1}$ ) containing  $I_n^{2n+1}$ , and, by Theorem 3.1,  $E[n]''$  has a subset  $X[n]''$  which is homeomorphic to  $X$ . Let

$$X[n] = \{x \in R^m \mid \exists y (\in X[n]'') \ x^{(l)} = y^{(l)} \ (l = 1, 2, \dots, 2n+1), \\ x^{(l)} = 0 \ (l = 2n+2, 2n+3, \dots, m)\}.$$

Then,  $X[n] \subset E[n]$ . Therefore

$$f'(X[n]) \subset C.$$

Therefore, since  $X$  is homeomorphic to  $X[n]$ ,  $X$  can be embedded in  $C$ .  $\square$

#### 4. $d(X) \leq \dim X$

**Theorem 4.1.** For any  $X (\in \mathcal{E})$ ,

$$d(X) \leq \dim X$$

(where  $d$  is an arbitrary function satisfying the conditions (A1)–(A5) in Section 1 for  $\mathcal{Q} = \mathcal{E}$ ).

**Proof.** It is obvious if  $\dim X = -1$ .

Let  $\dim X = n$  ( $0 \leq n < \infty$ ).

In  $R^{2n+1}$ ,  $I_n^{2n+1}$  is the sum of countably many  $n$ -parallelotopes. Since  $d(D) = n$  for any  $n$ -dimensional interval  $D$  from the conditions (A3) and (A5), we have  $d(I_n^{2n+1}) = n$  by condition (A2). By condition (A4), there exists a compactification  $A (\in \mathcal{E})$  of  $I_n^{2n+1}$  such that  $d(A) = d(I_n^{2n+1})$ . By Theorem 3.2,  $A$  has a subset  $X'$  which is homeomorphic to  $X$ . Therefore we have

$$d(X) = d(X') \leq d(A)$$

by conditions (A3) and (A1). Since

$$d(A) = d(I_n^{2n+1}) = n,$$

we have  $d(X) \leq n$ . And  $d(X) \leq \dim X$  since  $\dim X = n$ .  $\square$

#### 5. $d(X) = \dim X$

The proof of the following lemma is essentially due to Menger [7, p. 208].

**Lemma 5.1.** If  $X (\in \mathcal{E})$  has  $\dim X \geq 1$  and  $\mathcal{P} (\subset \mathcal{E})$  satisfies the following conditions (A1')–(A5'):

(A1') if  $Y' \subset Y$  and  $Y \in \mathcal{P}$ , then  $Y' \in \mathcal{P}$ ,

- (A2') if  $Y$  is the sum of countably many closed sets ( $\in \mathcal{P}$ ) of  $Y$ , then  $Y \in \mathcal{P}$ ,  
 (A3') if  $Y \in \mathcal{P}$  and  $Y$  is homeomorphic to  $Y'$ , then  $Y' \in \mathcal{P}$ ,  
 (A4') if  $Y \in \mathcal{P}$ , then there exists a compactification  $Y'$  of  $Y$  such that  $Y' \in \mathcal{P}$ ,  
 (A5')  $X \in \mathcal{P}$ ,

then  $\mathcal{P}$  contains a (1-dimensional) arc as an element.

**Proof.** (1) There exists a compactification  $X_0 (\in \mathcal{P})$  of  $X$ , and there exists a connected component  $X'$  of  $X_0$  containing at least two points. For some integer  $m (> 0)$ ,  $X' \subset I^m (\subset R^m)$ . Define the sets of  $m$ -cubes  $Q[n]$  ( $n = 0, 1, 2, \dots$ ) as follows:

Let  $Q[0] = \{I^m\}$ .

Divide  $I^m$  into  $2^{mn}$   $m$ -cubes each of which has edges of length  $(\frac{1}{2})^n$ , and let  $Q[n]$  be the set of these  $2^{mn}$   $m$ -cubes.

For each element  $Q(n, i)$  ( $i = 1, 2, \dots, mn$ ) of  $Q[n]$  ( $n = 0, 1, 2, \dots$ ), there exists a set  $X(n, i) (\subset Q(n, i))$  such that  $X(n, i)$  contains all vertices of  $Q(n, i)$  and is homeomorphic to  $X'$ . Let

$$Y = \bigcup \{X(n, i) \mid n = 0, 1, 2, \dots; i = 1, 2, \dots, mn\}.$$

Then  $Y \in \mathcal{P}$  since each  $X(n, i)$  is compact and is in  $\mathcal{P}$ . Obviously  $Y$  is connected and dense in  $I^m$ .

(2) Let  $S$  be an arbitrary set with  $Y \subset S \subset I^m$ . Then  $S$  is connected. We shall prove that  $S$  is locally connected.

Let  $p$  be an arbitrary point of  $S$  and  $U(p; \varepsilon)$  be the  $\varepsilon$ -neighborhood (in  $R^m$ ) of  $p$ , where  $\varepsilon$  is an arbitrary real number ( $> 0$ ). Let  $a, b$  ( $a \neq b$ ) be two arbitrary points of  $U(p; \varepsilon) \cap S$ . For some integer  $n$ , there exist finitely many elements  $Q(n, i_1), Q(n, i_2), \dots, Q(n, i_j)$  of  $Q[n]$  such that

$$\begin{aligned} a &\in Q(n, i_1), & b &\in Q(n, i_j), \\ Q(n, i_k) &\subset U(p; \varepsilon) \quad (k = 1, 2, \dots, j), \end{aligned}$$

and

$$Q(n, i_k) \cap Q(n, i_{k+1}) \neq \emptyset \quad (k = 1, 2, \dots, j-1).$$

Let  $P = \bigcup_{k=1}^j Q(n, i_k)$  and let

$$T = \bigcup \{X(n', i') \mid X(n', i') \subset P\}.$$

Then,  $T$  is connected and the closure of  $T$  is equal to  $P$ . Therefore  $T \cup \{a, b\}$  is connected, since  $T \subset T \cup \{a, b\} \subset P$ . It follows that  $U(p; \varepsilon) \cap S$  is connected. Thus  $S$  is locally connected.

(3) There exists a compactification  $Z (\in \mathcal{P})$  of  $Y$ . We may assume that  $Z \subset R^k$  ( $k \geq m$ ) for some integer  $k$ . Let  $Y'$  be the image of  $Y$  embedded in  $Z$  and  $f$  the embedding mapping  $Y \rightarrow Z$ . There exist two  $G_\delta$ -sets  $G, G'$  (of  $R^k$ ) and a homeomorphic mapping  $f': G \rightarrow G'$  such that  $Y \subset G$ ,  $Y' \subset G'$  and  $f'$  is an extension of  $f$ . Let

$$D = ((f')^{-1}(G' \cap Z)) \cap I^m (\subset R^k)$$

and let  $D' = f'(D)$ . Then both  $D$  and  $D'$  are  $G_\delta$ -sets (of  $R^k$ ). Since  $Z \in \mathcal{P}$ , we have  $D' \in \mathcal{P}$ , and therefore  $D \in \mathcal{P}$ . Since  $Y \subset D \subset I^m$ ,  $D$  is connected and locally connected by (2). Since  $D$  is a  $G_\delta$  in  $R^k$ , there exists a complete metric space  $C$  which is homeomorphic to  $D$  [2]. Since  $C$  is connected and locally connected,  $C$  has a (1-dimensional) arc as a subset. Therefore  $D$  has an arc as a subset, and it follows that  $\mathcal{P}$  has an arc as an element.  $\square$

**Lemma 5.2.** *If  $X (\in \mathcal{E})$  has  $\dim X \geq 1$ , then  $d(X) \geq 1$  (where  $d$  is an arbitrary set function satisfying the conditions (A1)–(A5) for  $\mathcal{Q} = \mathcal{E}$ ).*

**Proof.** Let  $d(X) = t$ . Since the class  $\mathcal{P}$  consisting of all  $X' (\in \mathcal{E})$  with  $d(X') \leq t$  satisfies the conditions (A1')–(A5') in Lemma 5.1,  $\mathcal{P}$  contains an arc  $L$  as an element. Since  $d(L) = 1$ , we have  $t \geq 1$ . Therefore  $d(X) \geq 1$ .  $\square$

**Theorem 5.3.** *For arbitrary  $X (\in \mathcal{E})$ ,*

$$d(X) \geq \dim X$$

*(where  $d$  is an arbitrary set function satisfying the conditions (A1)–(A6) for  $\mathcal{Q} = \mathcal{E}$ ).*

**Proof.** (1) If  $d(X) = -1$ , then  $\dim X = -1$ , therefore  $d(X) \geq \dim X$ .

(2) Let  $-1 < d(X) \leq 0$ . Since  $X$  is not empty,  $X$  has at least one point. Hence  $d(X) \geq 0$ , and therefore  $d(X) = 0$ . If we suppose  $\dim X \geq 1$ , then, by Lemma 5.2, we have  $d(X) \geq 1$ , and this contradicts the assumption  $d(X) \leq 0$ . Therefore  $d(X) \geq \dim X$ .

(3) Let  $d(X) = t$  ( $0 \leq n-1 \leq t < n$  where  $n$  is an integer). By (A6), there exist  $n$  subsets  $X_1, X_2, \dots, X_n$  of  $X$  such that

$$X = X_1 \cup X_2 \cup \dots \cup X_n, \quad d(X_i) \leq 0 \quad (i = 1, 2, \dots, n).$$

By (2),  $\dim X_i \leq 0$  ( $i = 1, 2, \dots, n$ ). Therefore

$$\dim X = \dim \left( \bigcup_{i=1}^n X_i \right) \leq n-1.$$

It follows that  $d(X) \geq \dim X$ .  $\square$

From Theorem 4.1 and Theorem 5.3, we have the following theorem:

**Theorem 5.4.** *For arbitrary  $X (\in \mathcal{E})$ ,*

$$d(X) = \dim X$$

*(where  $d$  is an arbitrary set function satisfying the conditions (A1)–(A6) for  $\mathcal{Q} = \mathcal{E}$ ).*

## 6. Independence of the axioms

We shall prove that the conditions (A1)–(A6) mentioned in Section 1 are independent for the case  $\mathcal{Q} = \mathcal{E}$ .



• *Independence of (A1).*

Let  $\mathcal{N}$  be the class of sets  $N$  such that  $N$  has  $\dim = 1$  and is a non-Borel subset of a Euclidean space. If  $X (\in \mathcal{E})$  has  $\dim = 1$  and satisfies the following condition  $\Delta$ :

$\Delta$ : for any countable closed covering  $\mathcal{C} = \{X_i | i = 1, 2, \dots\}$  of  $X$ , there exists a member  $X_i$  of  $\mathcal{C}$  such that  $X_i \in \mathcal{N}$ , then let  $d(X) = 2$ . And, for the other  $X (\in \mathcal{E})$ , let  $d(X) = \dim X$ .

(1) There exists  $X (\in \mathcal{E})$  having  $\dim = 1$  and  $d(X) = 2$ . (For example,  $X = N \times I$  ( $\subset R^2$ ) where  $N$  is a non-Borel set of  $I$ .) There exists a compactification  $X'$  of  $X$  such that  $\dim X' = \dim X = 1$ . Since  $X'$  is a Borel set in a Euclidean space and  $\dim X' = 1$ , we have  $d(X') = 1$ . As  $X$  is embedded in  $X'$ , this function  $d$  does not satisfy (A1).

(2) Let  $X \in \mathcal{E}$ ,  $X = \bigcup_{i=1}^{\infty} X_i$  and  $d(X_i) \leq 1$  ( $i = 1, 2, \dots$ ), where each  $X_i$  is a closed set of  $X$ . We shall prove  $d(X) \leq 1$ . Since  $d(X_i) \leq 1$ , for any countable closed covering  $\mathcal{C}_i = \{X_{i,j} | j = 1, 2, \dots\}$  of  $X_i$ , each  $X_{i,j}$  is either a non-Borel set with  $\dim = 0$  or a Borel set with  $\dim \leq 1$ . Since each  $X_{i,j}$  is closed in  $X$ , from the definition,  $d(X) \leq 1$ .

It is easily established that this function  $d$  satisfies (A2) for the other cases and also satisfies (A3)–(A6).

• *Independence of (A2).*

If  $X (\in \mathcal{E})$  is homeomorphic to a subset of the 1-dimensional interval  $I$ , then let  $d(X) = \dim X$ . If  $X (\in \mathcal{E})$  has  $\dim X = 1$  and  $X$  is not homeomorphic to any subset of  $I$ , then let  $d(X) = 2$ . And, for each space  $X (\in \mathcal{E})$  having  $\dim X \geq 2$ , let  $d(X) = \dim X$ .

• *Independence of (A3).*

Let  $\mathcal{B}$  be the class of sets  $B$  such that  $B$  has  $\dim = 1$  and is the image of a second category set of  $R^2$  by an isometric mapping. If  $X (\in \mathcal{E})$  has  $\dim = 1$  and has a metric subspace which is a member of  $\mathcal{B}$ , then let  $d(X) = 2$ . And, for the other  $X (\in \mathcal{E})$ , let  $d(X) = \dim X$ .

Let  $B$  be a second category subset of  $R^1$  with  $\dim B = 0$ , and  $C$  be a nowhere dense subset of  $R^1$  which is homeomorphic to  $B$ . Then,  $B \times I$  is homeomorphic to  $C \times I$ . However,  $d(B \times I) = 2$ ,  $d(C \times I) = 1$ . Therefore, this function  $d$  does not satisfy (A3). It is easily established that  $d$  satisfies (A1), (A2), (A4)–(A6).

• *Independence of (A4).*

Let  $\mathcal{B}$  be the class of sets having the form  $\{B | B \text{ is the sum of countably many closed subsets } B_1, B_2, \dots \text{ such that each } B_i \text{ is homeomorphic to a subset of the interval } I\}$ . If  $X (\in \mathcal{E})$  is a member of  $\mathcal{B}$  and has  $\dim X = 1$ , then let  $d(X) = 1$ . If  $X (\in \mathcal{E})$  is not a member of  $\mathcal{B}$  and has  $\dim X = 1$ , then let  $d(X) = 2$ . And, for the other  $X (\in \mathcal{E})$ , let  $d(X) = \dim X$ .

It is easily established that  $d$  satisfies (A1)–(A3), (A5), (A6). We shall prove that  $d$  does not satisfy (A4). Denote points of  $R^2$  by  $(x, y)$  where  $x$  and  $y$  are real

numbers. Let

$$C = \{(x, y) \in \mathbb{R}^2 \mid x \text{ is irrational, } 0 < x < 1, 0 \leq y \leq 1\}.$$

Then  $C$  has  $\dim = 1$ . But  $d(C) = 2$ , because, if  $C = \bigcup_{i=1}^{\infty} C_i$  where each  $C_i$  is closed in  $C$ , then some  $C_i$  contains uncountably many pairwise disjoint segments. Therefore  $C_i \not\subset I$ , hence  $C \notin \mathcal{B}$ . On the other hand,  $d(I_1^3) = 1$ , since  $I_1^3$  is the sum of countably many segments. By Theorem 3.2, any compactification  $A$  of  $I_1^3$  contains a subset which is homeomorphic to  $C$ . Therefore  $d(X) \geq 2$ . Therefore, there is no compactification  $A$  of  $I_1^3$  such that  $d(I_1^3) = d(A)$ . Hence, this function  $d$  does not satisfy (A4).

• *Independence of (A5).*

Let  $d(\emptyset) = -1$ , and, let  $d(X) = 0$  for every  $X \in \mathcal{E}$  which is not empty.

• *Independence of (A6).*

Let  $Z(p)$  be the cyclic group of order  $p$ , where  $p$  is a prime number. If we let  $d(X) = \text{c-dim}_{Z(p)} X$  where  $\text{c-dim}_{Z(p)} X$  is the cohomological dimension of  $X \in \mathcal{E}$ , then  $d(X)$  satisfies the conditions (A1)–(A5) [4] but does not satisfy the condition (A6). (Because, if we suppose that  $d(X)$  satisfies the conditions (A1)–(A6), then, by Theorem 5.4, we have  $d(X) = \dim X$  for every  $X \in \mathcal{E}$ , and is contradictory to  $\text{c-dim}_{Z(p)} X \neq \dim X$  for some  $X$ .)

## References

- [1] T.M. Aarts, A characterization of strong inductive dimension, *Fund. Math.* 70 (1971) 147–155.
- [2] P. Alexandroff, Sur les ensembles de la première classe et les espaces abstraits, *C. R. Acad. Sci. Paris* 178 (1924) 185–187.
- [3] P.S. Alexandrov, The axiomatic construction of dimension theory, in: *Proceedings of the International Symposium on Topology and its Applications, Budva (1972)* 16–17 (in Russian).
- [4] V.I. Kuz'minov, Homological dimension theory, *Uspekhi Mat. Nauk* 23 (5) (1968) 3–50 (in Russian); also: *Russian Math. Surveys* 23 (5) (1968) 1–45 (in English).
- [5] M.M. Lavrentieff, Contribution à la théorie des ensembles homéomorphe, *Fund. Math.* 6 (1924) 149–160.
- [6] O.V. Lokucievskii and E.V. Ščepin, On P.S. Alexandroff's axiomatic definition of the dimension, topics in topology, in: *Proceedings Colloquium, Keszthely (1972)* 477–480; also: *Colloquia Mathematica Societatis János Bolyai* 8 (North-Holland, Amsterdam, 1974).
- [7] K. Menger, Zur Begründung einer axiomatischen Theorie der Dimension, *Monatsh. Math.* 36 (1929) 193–218.
- [8] T. Nishiura, Inductive invariants and dimension theory, *Fund. Math.* 59 (1966) 243–262.
- [9] T. Nishiura, An axiomatic characterization of covering dimension in metrizable spaces, in: *Proceedings Second Pittsburgh International Conference, Carnegie-Mellon University and University of Pittsburgh, PA (1972)* 341–353.
- [10] S. Sakai, An axiomatic characterization of the large inductive dimension for metric spaces, *Proc. Japan Acad.* 44 (1968) 782–785.
- [11] E. Ščepin, An axiomatization of the dimension of metric spaces, *Dokl. Akad. Nauk SSSR* 206 (1972) 31–32 (in Russian); also: *Soviet Math. Dokl.* 13 (1972) 1177–1179 (in English).
- [12] E.V. Ščepin, An axiomatization of the dimension of metric spaces, *Mat. Sb. (N.S.)* 92 (1973) 135–141 (in Russian); also: *Math. USSR-Sb.* 21 (1973) 137–143 (in English).
- [13] I.G. Tsereteli, Axiomatic characteristics of dimension dim on special classes of spaces and their continuous mappings, *Soobshch. Akad. Nauk Gruz. SSR* 122 (1) (1986) 29–32.